


regeneration

Regeneration of the anterior mandible: A clinical case presentation



John Lupovici, DDS

Although autogenous bone has long been considered to be the gold standard for restoring deficient alveolar bone, a number of drawbacks have been associated with its use. Alternative grafting materials have developed, along with techniques for maintaining existing soft tissue and bone. This article briefly reviews these developments. A clinical treatment is also presented in which a new xenograft material was used in combination with a cross-linked, cell-occlusive membrane to restore a severely resorbed edentulous mandible in preparation for placement of dental implants.

Key Words: regeneration, bone grafting, grafting materials, xenograft

Introduction

An estimated 1,260,000 dental bone-grafting procedures were performed in the United States in 2006, and that number is expected to grow by more than 15% annually.¹ As hard-tissue reconstruction has become an increasingly routine part of dental surgical care, demand for suitable grafting materials has increased. Such materials must satisfy various regulatory requirements and meet clinicians' expectations for safety and effectiveness. Ideally, they should be biocompatible, easy to procure, resorbable, osseointegrative, osseoconductive, and cost-effective.

In response to these considerations, autografts, allografts, alloplasts, and xenografts have all become acceptable alternatives for filling and regenerating bone defects. However, autograft material may be difficult to obtain in sufficient quantity, and its harvest poses risks of pain, complications, and morbidity.²⁻⁷ Moreover, high resorption rates and limited viability have been reported.⁸⁻¹¹

Allograft bone, while more easily obtainable, may lack osseointegrativity (depending upon its source and processing).¹²⁻¹⁴ Alloplasts typically lack osseointegrativity, and may have variable resorption rates.

Xenografts constitute the fourth category of commonly used bone-grafting materials, with porous bovine-derived material—the most popular xenograft variety. This biocompatible material may eliminate the need for a second surgical site and may serve as an effective regenerative matrix for a variety of indications prior to implant placement. Numerous researchers have reported a high degree of osseointegrativity,¹⁵⁻²² and bovine bone particles are well incorporated within newly regenerated grafted bone, according to histological findings. It has been argued that the slow resorption profile of bovine-derived bone may contribute to increased stability of the regenerated bone.²³

Traditionally most xenograft used in oral regenerative procedures has been unsintered. Such material undergoes a multi-step process of annealing (up to 300 degrees Celcius), followed by treatment with organic solvents such as sodium hydroxide.²⁴ However, a second type of bovine-derived bone, used since 1989 as a bone-replacement material in the fields of orthopedic and skull surgery, has recently been introduced into the field of oral regeneration. This material uses a process known as sintering to remove all pathogenic components and organic components from the bovine bone. The bovine bone is heated to more than 1200 degrees Celcius, yielding a highly crystalline material containing small amounts of calcium oxide resulting from decomposition of the original carbon content. The sintered bovine bone incorporates its native macroporosities, as well as preserves its microposity of the original bone.

The following clinical case presentation illustrates the use of sintered xenograft particles in combination with a resorbable collagen membrane to regenerate an edentulous mandible.

The patient was a 72-year-old female who presented with an ill-fitting denture following long-standing mandibular edentulism. Clinical and radiographic examination revealed a narrow alveolar ridge with significant apical undercuts (Figs. 1 and 2). The patient required pre-operative ridge regeneration to accommodate a future implant-supported overdenture.

Sintered bovine bone particles were chosen as the graft material due to their biologic and physical properties. Biologic properties include the ability to function as an osteoconductive regenerative material. The physical characteristics of bovine bone offer the additional advantage of increased compressive resistance, due to the material's inherent native structure. Regenerative attempts with other graft materials may be compromised by their inability to contend with the compressive forces delivered by an overlying denture.

Following treatment-plan acceptance, the patient was anesthetized, and a full-thickness mucoperiosteal flap was reflected (Fig. 3). Upon identification of the mental foramina, distal releasing incisions were made. Measurement of the existing ridge indicated that the width was 2-3mm wide (Fig. 4).



Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5



Fig. 6

Fig. 7



Fig. 8




Fig. 9



Fig.10



Fig.11





Fig.12



The residual periosteum was carefully debrided with hand instruments, and the ridge was decorticated using a high speed round carbide bur with copious irrigation (Fig. 5).

After hydration in sterile saline for five minutes, Endobon® Xenograft Granules (BIOMET **3i**) were molded over the decorticated ridge. Excess graft material was applied to allow for the characteristic shrinkage associated with guided bone regeneration and the unavoidable compression by the denture (Figs. 6 and 7).

Graft containment was achieved by tucking cross-linked OsseoGuard® Resorbable Collagen Membranes (BIOMET **3i**) under the facial flap and draping them over the graft (Fig. 8). Extensive periosteal releasing incisions of the buccal flap in conjunction with the initial lingual flap reflection to the floor of the mouth enabled passive primary closure. Continuous locking 4.0 Gore-Tex® Sutures (W.L. Gore & Associates, Flagstaff, Arizona, USA) secured the soft-tissue flaps (Fig. 9).

The intaglio surface of the existing denture was relieved and relined to reduce the potential compressive forces on the regenerated area, as well as to maintain primary closure. The patient was then released with analgesics and antibiotics.

Post-operative healing was uneventful, and primary closure of the regenerated site was maintained throughout the healing period. Six months later, the patient was seen for evaluation and placement of dental implants (Fig. 10). A crestal incision was made, and full-thickness periosteal flaps were reflected to reveal a sufficient gain in ridge width to support implant placement (Fig. 11).

Following the manufacturer's recommended protocol, osteotomies were performed in the two cuspid sites for placement of 4mm diameter NanoTite™ Certain® Implants (BIOMET **3i**) (Fig. 12). Cover screws were placed into the internal interfaces of the implants and tightened by hand. A 2mm trephine core was harvested from the central incisor location, and histological evaluation revealed excellent incorporation of the Endobon Xenograft Granules. The soft-tissue flaps were closed with continuous locking sutures, and the intaglio surface of the existing denture was relieved and relined. The patient was then released with analgesics and antibiotics.

Three months later, after uneventful healing, a tissue punch was used to expose the implants. The cover screws were removed, and EP® Healing Abutments (BIOMET 3i) were placed. The denture was relieved over the healing abutments, and the patient was dismissed with oral hygiene instructions.

Following soft-tissue maturation at eight weeks, the patient was seen by the restorative clinician for impressions and fabrication of the definitive prosthesis. The mandibular overdenture will be retained by Locator® Abutments (BIOMET 3i) processed directly into the overdenture base.

Clinical Relevance

For many severely resorbed edentulous patients, being unable to wear a removable denture during healing may be a significant impediment to accepting treatment with dental implants. Treatment plans that acknowledge this psychological reality are likely to enjoy higher acceptance rates. Sintered bovine-derived bone particles used in combination with cross-linked, cell-occlusive membranes may enable clinicians to obtain successful regenerative outcomes in cases where compressive forces imposed by dentures might otherwise compromise the success of the augmentation or lead to questionable regenerative gain.

References

- Millennium Research Group, US Market Dental Biomaterials Survey 2007 Toronto, Canada.
- Misch C. Comparison of intraoral donor sites for onlay grafting prior to implant placement. *Int J Oral Maxillofac Implants* 1997;12:767-776.
- Pikos MA. Block autografts for localized ridge augmentation: Part I. The posterior maxilla. *Implant Dent* 1999;8:279-285.
- Rudman R. Prospective evaluation of morbidity associated with iliac crest harvest for alveolar cleft grafting. *J Oral Maxillofac Surg* 1997;55:2219-2223.
- Schwartz-Arad D, Levin L, Sigal L. Surgical success of intraoral autogenous block onlay grafting for alveolar ridge augmentation. *Implant Dent* 2005;14:131-138.
- Raghoobar G, Louwervse C, Kalk W et al. Morbidity of chin bone harvesting. *Clin Oral Implants Res* 2001;12:503-507.
- Clavero J, Lundgren S. Ramus or chin grafts for maxillary sinus inlay and local onlay augmentation: comparison of donor site morbidity and complications. *Clin Implant Dent Relat Res* 2003;5:215-218.
- Misch C, Misch C, Resnik R et al. Reconstruction of maxillary alveolar defects with mandibular symphysis grafts for dental implants: A preliminary procedural report. *Int J Oral Maxillofac Implants* 1992;7:360-366.
- Widmark G, Andersson B, Ivanoff C. Mandibular bone graft in the anterior maxilla for single-tooth implants. Presentation of surgical method. *Int J Oral Maxillofac Surg* 1997;26:106-109.
- McAllister B, Haghight K. Bone augmentation techniques. *J Periodontol* 2007;78:377-396.
- Proussaefs P, Lozada J. The use of intraorally harvested autogenous block grafts for vertical alveolar ridge augmentation: A human study. *Int J Periodontics Restorative Dent* 2005;25:351-363.
- Becker W, Becker B, Caffesse R. A comparison of demineralized freeze-dried bone and autogenous bone to induce bone formation in human extraction sockets. *J Periodontol* 1994;65:1128-1133.
- Becker W, Clokie C, Sennerby L et al. Histologic findings after implantation and evaluation of different graft materials and titanium micro screws into extraction sockets: Case report. *J Periodontol* 1998;69:414-421.
- Schwartz Z, Sommers A, Mellonig J et al. Ability of commercially demineralized freeze-dried bone allograft to induce new bone formation. *J Periodontol* 1996;68:1085-1092.
- Artzi Z, Tal H, Dayan D. Porous bovine bone mineral in healing of human extraction sockets: 2. Histochemical observations at 9 months. *J Periodontol* 2001;72:152-159.
- Peetz M. Characterization of xenogenic bone material. In: Boyne PJ, Evensen L, Eds. *Osseous reconstruction of the maxilla and the mandible: Surgical techniques using titanium mesh and bone mineral*. Carol Stream, IL: Quintessence 1997:87-100.
- McAllister BS, Margolin MD, Cogan AG et al. Residual lateral wall defects following sinus grafting with recombinant human osteogenic protein-1 or Bio-Oss in the chimpanzee. *Int J Periodontics Restorative Dent* 1998;18:227-239.
- Klinge B, Alberius P, Isaksson S et al. Osseous response to implanted natural bone mineral and synthetic hydroxylapatite ceramics in the repair of experimental skull bone defects. *J Oral Maxillofac Surg* 1992;50:241-249.
- Spector M. Anorganic bovine bone and ceramic analogs of bone mineral as implants to facilitate bone regeneration. *Clin Plast Surg* 1994;21:437-444.
- Hammerle CHF, Alah AJ, Schmid J et al. The biological effect of deproteinized bovine bone on bone neof ormation on the rabbit skull. *Clin Oral Implants Res* 1997;8:198-207.
- Hammerle CHF, Chiantella Gc, Karring T et al. The effect of a deproteinized bovine bone mineral on bone regeneration around titanium dental implants. *Clin Oral Implants Res* 1998;9:151-162.
- Jensen SS, Aaboe M, Pinholt EM et al. Tissue reaction and material characteristics of four bone substitutes. *Int J Oral Maxillofac Implants* 1996;11:55-66.
- Simion M, Jovanovic SA, Tinti et al. Long-term evaluation of osseointegrated implants inserted at the time or after vertical ridge augmentation. A retrospective study on 123 implants with 1-5 year follow-up. *Clin Oral Implants Res* 2001;12(1):35-45.
- Tadic D, Epple M. A thorough physicochemical characterization of 14 calcium phosphate-based bone substitution materials in comparison to natural bone. *Biomaterials* 2004;25:987-994.

John Lupovici, DDS



Dr. Lupovici received his dental degree and certificate in Periodontology from New York University. While at NYU, he participated in numerous research studies, which resulted in his being awarded three Dean's Student Research Awards and receiving first place in the American Dental Association / NYU Research Day Competition. He is a Diplomate of the American Board of Periodontology and holds a faculty position at NYU in the Department of Periodontics and Implant Dentistry. His clinical research includes such topics as bone regeneration and implant dentistry; subjects on which he has published and lectured nationally and internationally. Dr. Lupovici maintains a private periodontal practice in New York City and Commack, New York.